

Detection of potential green open space area using landsat 8 satellite imagery

by Aditya Galih Sulaksono

Submission date: 12-Jun-2023 10:58AM (UTC+0700)

Submission ID: 2114157286

File name: Detection_of_potential_green_open_space.pdf (676.15K)

Word count: 3439

Character count: 18574



Research paper

doi: 10.30822/arteks.v6i1.730

Detection of potential green open space area using landsat 8 satellite imagery

Respati Wikantiyoso^{1*}, Aditya Galih Sulaksono², Tonny Suhartono¹

¹Department of Architecture, Universitas Merdeka Malang

²Department of Information System, Universitas Merdeka Malang

Terusan Raya Dieng, 62-64, Malang, Indonesia



ARTICLE INFO	ABSTRACT
<p><i>Article history:</i> Received November 04, 2020 Received in revised form November 26, 2020 Accepted November 28, 2020 Available online April 01, 2020</p> <p><i>Keywords:</i> Green open space Landsat 8 NDVI</p> <p>*Corresponding author: Respati Wikantiyoso Department of Architecture Universitas Merdeka Malang, Indonesia Email: adit@unmer.ac.id ORCID: https://orcid.org/0000-0003-4301-7868</p>	<p><i>Green city development policies have an essential role in improving the ecological functions of a municipality. According to Law 26/2007 on spatial planning, it contains provisions for green open space which oblige the government to set aside 30% of the city as green open space. However, the main obstacle in providing green open space remains the limited availability of urban land. Therefore, efforts are needed to detect urban land that is available for use as green open space. The purpose of this study is to determine the initial description of the availability of green open space in Malang City. This study uses an integrated method of calculating the Normalized Difference Vegetation Index (NDVI) with land cover classification. NDVI analysis reveals the distribution value of vegetation density in Malang City. Based on NDVI analysis results, green open space requires medium and high density vegetation. In Malang City, only 2946.56 ha, or 26.60% of the city, meets these requirements. The availability of green open space in Malang City therefore does not meet minimum green open space requirements. This study's results suggest the need for additional policies aimed at providing green open space for Malang City.</i></p>

Introduction

Green open space (GOS), as an urban landscape element, exhibits a city's social, cultural, economic, and ecological dimensions. Thus, urban GOS has a very strategic function to create an urban environment that promotes harmonious and sustainable interaction between human life and the natural environment (Respati Wikantiyoso et al. 2020). The Minister of Home Affairs Regulation No. 1 of 2007 concerns the preparation of urban GOS. According to this regulation, urban GOS planning ideally requires an area of green space equal to at least 30% of the city (R Wikantiyoso, Suhartono, and Sulaksono 2020). This 30% should be comprised of 20% public GOS and 10% private GOS in each region.

Providing and utilizing a city's open space to achieve a 30% balance is a complicated problem for local governments in urban development (Respati et al. 2020).

The main obstacle in providing green open space is the limited availability of urban land. Thus, efforts are needed to detect urban land that is available for use as green open space. Efforts to realize the conceptual provision of urban open space planning have produced regulatory requirements related to the provision of urban public spaces (Juwito, Wikantiyoso, and Tutuko 2019). Still, technical regulations cannot function as guidelines and directives for improving the quality of open urban areas (Pannell and Roberts 2009).



Vegetation is an important constituent of land use, and it exhibits diverse types (Ives et al. 2014). The collection of various vegetation types will produce different vegetation density levels, and these levels, in turn, impact land use in an area. Current efforts aim to use technology to assess the level of vegetation density in an area (Ives et al. 2014; Lausch et al. 2018; López et al. 2001). Vegetation has unique spectral characteristics that allow analysts to obtain an index representing the vegetation's condition using various methods. These technologies are remote sensing technology and geographic information systems (GIS).

The method of measuring vegetation using satellite imagery relies on the reflectance of landscape features (Mears and Brindley 2019). Vegetation index results from remote sensing data processing can note the presence and condition of vegetation in urban areas (Yunhao et al. 2006). The data utilization of remote sensing satellites for green open space mapping offers numerous advantages over the terrestrial method (López et al. 2001; Li, Fan, and Shen 2018; Kothencz et al. 2017; Mears and Brindley 2019). The method commonly used for GOS mapping employs satellite data for qualitative classification, or quantitative classification using a vegetation index. Meanwhile, green open space mapping research uses Normalize Difference Vegetation Index (NDVI) to measure plant growth and determine the vegetation coverage area (Mears and Brindley 2019; Mohamed, Worku, and Kindu 2019; Aguilera Benavente, Rodríguez Espinosa, and Gómez Delgado 2018). In NDVI calculations, the wavelength of visible red and near-infrared (NIR) light is used. The basic calculation is based on the pigment inside the leaves or chlorophyll highly absorbing light (0.4 – 0.7 μm) during the photosynthesis process. Meanwhile, the leaves' cell structure is highly reflective of near-infrared light (0.7 – 1.1 μm) (Sulma et al. 2017).

This research was conducted to analyze the vegetation density level of green open space in Malang using Landsat 8 satellite imagery data with the NDVI method. NDVI is the standard method for comparing the level of the greenness of vegetation in satellite image data. NDVI can be used as an indicator of biomass, relative greenness, and the status (in terms of health and density) of vegetation in an area (Lausch et al. 2018; Lai et al. 2019; Li, Fan, and Shen 2018). This study aimed to describe the potential availability of green open space in Malang using

an integrated NDVI calculation method with Landsat 8 OLI data.

Table 1. Landsat 8 bands designation

Bands	Resolution (m)	Wavelength (μm)
Band 1 – Coastal aerosol	30	0.435 – 0.451
Band 2 – Blue	30	0.452 – 0.512
Band 3 – Green	30	0.533 – 0.590
Band 4 – Red	30	0.636 – 0.673
Band 5 – NIR	30	0.851 – 0.879
Band 6 – SWIR 1	30	1.566 – 1.651
Band 7 – SWIR 2	30	2.107 – 2.294
Band 8 – Panchromatic	15	0.503 – 0.676
Band 9 – Cirrus	30	1.363 – 1.384
Band 10 – TIRS1	100	10.60 – 11.19
Band 11 – TIRS2	100	11.50 – 12.51

Source: NASA Landsat 8 Bands

Landsat 8 launched on February 11, 2013. It was originally called the Landsat Data Continuity Mission (LDCM), and it carried two sensors: An On-board Operational Land Imagery (OLI) sensor and a Thermal Infrared Sensor (TIRS) with 11 channels. Among these channels, nine channels (bands 1-9) are on OLI, while two others (bands 10 and 11) are on TIRS. Table 1 shows the channel characteristics of Landsat 8 satellite imagery.

Method

Research on the potential availability of green open space in Malang uses satellite image data from Landsat 8 OLI Level 1TP sensor line 118, line 066. Figure 1 show the recording from September 17, 2020, which was obtained from the LAPAN Landsat Catalog. This research used the descriptive interpretation method with NDVI analysis of Landsat 8 OLI imagery.




Figure 1. Landsat 8 satellite imagery path/row 118/066
Source: LAPAN Landsat Catalog, 2020-09-17

The research began with preprocessing the Landsat 8 satellite image data by making geometric corrections and cropping the image to create a subset image. Landsat 8 image data was then used for further analysis (A. Purwanto 2015; Xu 2014). This research used the NDVI calculation method with land cover classification integration (Lausch et al. 2018; Li, Fan, and Shen 2018).

NDVI analysis produces the distribution value of vegetation density in Malang City. NDVI is sensitive to chlorophyll's photosynthetic activity, which enables researchers to use the NDVI value to classify vegetation (Zaitunah et al. 2018; A. D. Purwanto et al. 2014). The number and thickness of a plant's leaves will significantly affect the results of its reflection (Lufilah, Makalew, and Sulistyantara 2017). If NIR reflection is greater than red wavelength radiation, an area's vegetation is dense, and it may be classified as forest. If there is a small difference between the brightness of the reflected red and NIR wavelengths, the vegetation can be classified as grasslands or rice fields in terms of vegetative mass (Ridwan and Muharoroh 2017). NDVI values represent vegetation classification based on plant dominance (A. D. Purwanto et al. 2014; Zaitunah et al. 2018).

A faster and more accurate calculation of green open space's distribution and potential can be conducted using the NDVI analysis method. The method and use of the latest Landsat 8 data can identify changes, including additions and reductions (Feltynowski et al. 2018; Rafiee,

Salman Mahiny, and Khorasani 2009; A. D. Purwanto et al. 2014), of green open space potential in Malang City. This method is an efficient calculation and can form the basis for policy decisions on Green Open Space provision for the city in the future.

Normalized Difference Vegetation Index (NDVI)

Data analysis in this study used the NDVI method according to the following equation:

$$NDVI = \frac{NIR - Red}{NIR + Red}$$

NIR wavelengths show from band five and red from band four in Landsat 8 satellite imagery data. NDVI values range from -1 to 1. NDVI values below 0 represent the presence of clouds, ice, or snow, while values between 0 and 0.1 usually indicate empty land, and values above 0.1 indicate vegetation (Febrianti and Sofan 2014).

Result and discussion

Based on NDVI analysis, the distribution of vegetation density values in Malang City ranged from -0.0078 to 0.5169 (figure 2). The level of vegetation density based on the NDVI value can be used as the basis for classification according to plant dominance (A. D. Purwanto et al. 2014). Vegetation surfaces with an NDVI value around 0.1 indicate grasslands and shrubs. Meanwhile, values between 0.1 and 0.8 indicate tropical rain forests, and NDVI values close to +1 indicate vegetation cover.




Figure 2. Malang NDVI
Source: LAPAN Landsat Catalog, 2020-09-17

In this study, the vegetation threshold value is 0.1836. A value less than the threshold value is a non-vegetation area occupied by buildings or other types of land (figure 3). The color spectrum in figure 3 shows the distribution of non-vegetation areas in Malang City. The color spectrum's identification with an NDVI index between -0.0078 to 0.1835 classifies it as a non-vegetation area (see table 1).



Figure 3. Non-vegetation area

Table 2 shows the results of the NDVI classification for Malang City in 2020. The non-vegetation class falls in the range of 0.0078 to 0.1835 and covers 5185.71 ha, or 46.81% of the total area of the city. The moderate vegetation class, which mostly includes shrubs and tall grasses, falls in the range of 0.1836 to 0.2506. It covers 29.59% of the total vegetation area, or 3278.16 ha. Finally, dense vegetation classes, like mangrove forests, have medium to dense canopies and are considered city forests. They fall in the range of 0.5207 to 0.5169 and comprise 23.60% of the total vegetation area, or around 2614.95 ha.

Figure 4 shows the green color spectrum distribution. The distribution and value of NDVI indicate the potential area and area of Green Open Space area. Based on the NDVI calculation of the area of vegetation (Zaitunah et al. 2018) in Malang City (figure 4), the potential of green open space in Malang City is 26.60%, which falls short of the 30% required under the Minister of Home Affairs Regulation No. 1 of 2007.



Figure 4. Vegetation area

Table 2. NDVI value of Malang City

No	Classification	NDVI value	Area (ha)	%
1	Non-vegetation	-0.0078 – 0.1835	5185.71	46.81
2	Moderate vegetation	0.1836 – 0.2506	3278.16	29.59
3	Dense vegetation	0.5207 – 0.5169	2614.95	23.60
Total area			11078.82	100

Table 3 shows the results of the NDVI analysis of vegetation density in each district of Malang City, including districts with the lowest vegetation densities and the highest vegetation densities. The district with the highest vegetation density was the Kedungkandang district at 19.03%, while the district with the lowest vegetation density was the Klojen district at 2.00%.

Table 3. Vegetation area by district of Malang City

No	District	NDVI value	Area (ha)	%
1	Kedungkandang	0.25907 – 0.516138	2108.43	19.03
2	Klojen	0.17511 – 0.450339	221.31	2.00
3	Blimbing	0.211059 – 0.516935	467.82	4.22
4	Lowokwaru	0.217884 – 0.505209	732.87	5.52
5	Sukun	0.216061 – 0.480683	722.88	6.52

Conclusion

The NDVI value distribution method offers a faster and more accurate calculation of the distribution and potential of green open spaces.

The method and use of the latest Landsat 8 data can identify changes, including additions and subtractions, of potential green open space in Malang City. Because it is more efficient while remaining accurate, this method may be useful as the government works to implement future green open space policies.

The novelty of research with this method lies in implementing a control mechanism for urban green open space availability, which requires 30% of the city area. Controlling changes in urban green open space availability is significant for controlling urban green open space provision. The detection method for green open space potential is significant and should be integrated with Green Open Space Management Information System development to highlight the potential for integrated and sustainable control of urban green open space provisions.

This research opens the opportunity for further detailed research by calculating the potential of green open space in river corridors, high-voltage electricity connection corridors and railway corridors, with more comprehensive parameters.

Acknowledgment

This paper is one of the research outcomes from Web-based Green City MIS as a Model for Implementing Sustainable Urban GOS Control. This research was funded by a Higher Education Leading Applied Research Grant from the Directorate of Research and Community Service, Ministry of Education and Culture of Indonesia. We thank the government of Malang City for permitting us to use spatial data in this research.

References

- Aguilera Benavente, Francisco, Víctor Manuel Rodríguez Espinosa, and Montserrat Gómez Delgado. 2018. 'Definición de Infraestructuras Verdes: Una Propuesta Metodológica Integrada Mediante Análisis Espacial'. *Documents d'Anàlisi Geogràfica* 64 (2): 313. <https://doi.org/10.5565/rev/dag.419>.
- Febrianti, Nur, and Parwati Sofan. 2014. 'Ruang Terbuka Hijau Di Dki Jakarta Berdasarkan Analisis Spasial Dan Spektral Data Landsat 8'. *Deteksi Parameter Geobiofisik Dan Diseminasi Penginderaan Jauh , Seminar Nasional Penginderaan Jauh*, no. April: 498–504.
- Feltynowski, Marcin, Jakub Kronenberg, Tomasz Bergier, Nadja Kabisch, Edyta Łaszkiewicz, and Michael W. Strohbach. 2018. 'Challenges of Urban Green Space Management in the Face of Using Inadequate Data'. *Urban Forestry & Urban Greening* 31 (April): 56–66. <https://doi.org/10.1016/j.ufug.2017.12.003>.
- Ives, Christopher D., Cathy Oke, Benjamin Cooke, Ascelin Gordon, and Sarah Bekessy. 2014. 'Planning for Green Open Space in Urbanising Landscapes'. Australia.
- Juwito, Juwito, Respati Wikantiyoso, and Pindo Tutuko. 2019. 'Kajian Persentase Ruang Terbuka Hijau Pada Implementasi Revitalisasi Taman Kota Malang'. *Local Wisdom : Jurnal Ilmiah Kajian Kearifan Lokal* 11 (1). <https://doi.org/10.26905/lw.v11i1.2686>.
- Kothencz, Gyula, Ronald Kolcsár, Pablo Cabrera-Barona, and Péter Szilassi. 2017. 'Urban Green Space Perception and Its Contribution to Well-Being'. *International Journal of Environmental Research and Public Health* 14 (7): 766. <https://doi.org/10.3390/ijerph14070766>.
- Lai, Hakkan, Emily J Flies, Philip Weinstein, and Alistair Woodward. 2019. 'The Impact of Green Space and Biodiversity on Health'. *Frontiers in Ecology and the Environment* 17 (7): 383–90. <https://doi.org/10.1002/fee.2077>.
- Lausch, Angela, Olaf Bastian, Stefan Klotz, Pedro J. Leitão, Andrés Jung, Duccio Rocchini, Michael E. Schaepman, Andrew K. Skidmore, Lutz Tischendorf, and Sonja Knapp. 2018. 'Understanding and Assessing Vegetation Health by in Situ Species and Remote-sensing Approaches'. Edited by Petteri Vihervaara. *Methods in Ecology and Evolution* 9 (8): 1799–1809. <https://doi.org/10.1111/2041-210X.13025>.
- Li, Zhiming, Zhengxi Fan, and Shiguang Shen. 2018. 'Urban Green Space Suitability Evaluation Based on the AHP-CV Combined Weight Method: A Case Study of Fuping County, China'. *Sustainability* 10 (8): 2656. <https://doi.org/10.3390/su10082656>.
- López, Erna, Gerardo Bocco, Manuel Mendoza, and Emilio Duhau. 2001. 'Predicting Land-Cover and Land-Use Change in the Urban Fringe'. *Landscape and Urban Planning* 55

- 12
(4): 271–85. [https://doi.org/10.1016/S0169-2046\(01\)00160-8](https://doi.org/10.1016/S0169-2046(01)00160-8).
- Lufilah, Siti Novianti, Afra DN Makalew, and Bambang Sulistyantara. 2017. 'Pemanfaatan Citra Landsat 8 Untuk Analisis Indeks Vegetasi Di DKI Jakarta'. *Jurnal Lanskap Indonesia*, 73–80. <https://doi.org/10.29244/jli.2017.9.1.73-80>.
- Meais, Meghann, and Paul Brindley. 2019. 'Measuring Urban Greenspace Distribution Equity: The Importance of Appropriate Methodological Approaches'. *ISPRS International Journal of Geo-Information* 8 (6): 286. <https://doi.org/10.3390/ijgi8060286>.
- Mohamed, Asfaw, Hailu Worku, and Mengistie Kindu. 2019. 'Quantification and Mapping of the Spatial Landscape Pattern and Its Planning and Management Implications a Case Study in Addis Ababa and the Surrounding Area, Ethiopia'. *Geology, Ecology, and Landscapes*, December, 1–12.
- Pannell, David J., and Anna M. Roberts. 2009. 'Conducting and Delivering Integrated Research to Influence Land-Use Policy: Salinity Policy in Australia'. *Environmental Science & Policy* 12 (8): 1088–98. <https://doi.org/10.1016/j.envsci.2008.12.005>.
- Purwanto, Ajun. 2015. 'Pemanfaatan Citra Landsat 8 Untuk Identifikasi Normalized Difference Vegetation Index (NDVI) Di Kecamatan Silat Hilir Kabupaten Kapuas Hulu'. *EDUKASI: Jurnal Pendidikan (e-Journal)* 13 (1): 27–36. <https://doi.org/http://dx.doi.org/10.31571/edukasi.v13i1.17>.
- 6
Purwanto, Anang Dwi, Wikanti Asriningrum, Gathot Winarso, and Ety Parwati. 2014. 'Analisis Sebaran Dan Kerapatan Mangrove Menggunakan Citra Landsat 8 Di Segara Anakan, Cilacap'. In *Seminar Nasional Penginderaan Jauh 2014: Pengolahan Data Dan Pengenalan Pola*, 232–41.
- Rafiee, Reza, Abdolrassoul Salman Mahiny, and Nematollah Khorasani. 2009. 'Assessment of Changes in Urban Green Spaces of Mashad City Using Satellite Data'. *International Journal of Applied Earth Observation and Geoinformation* 11 (6): 431–38. <https://doi.org/10.1016/j.jag.2009.08.005>.
- Respati, Wikantiyoso, Suhartono Tonny, Sulaksono Aditya Galih, and Wikananda Triska Prakasa. 2020. 'Corporate Social Responsibility (CSR) Model in Improving the Quality of Green Open Space (GOS) to Create a Livable City'. In *Corporate Social Responsibility [Working Title]*. IntechOpen. <https://doi.org/10.5772/intechopen.94481>.
- Ridwan, Martin, and Evi Muharoroh. 2017. 'Pemanfaatan Penginderaan Jauh Dalam Perhitungan Luas Ruang Terbuka Hijau (RTH): Studi Kasus Kota Bekasi'. In *Indonesian Science Student Conference 2017*, 16–21. Yogyakarta.
- Sulma, Sayidah, Jalu Tejo Nugroho, Any Zubaidah, Hana Listi Fitriana, and Nanik Suryo Haryani. 2017. 'Detection of Green Open Space Using Combination Index of Landsat 8 Data (Case Study: Dki Jakarta)'. *International Journal of Remote Sensing and Earth Sciences (IJReSES)* 13 (1): 1.
- Wikantiyoso, R, T Suhartono, and A G Sulaksono. 2020. 'Controlling Efforts of Green Open Space Provision in East Malang Residential Areas Development, Indonesia'. *IOP Conference Series: Earth and Environmental Science* 562 (September): 012015. <https://doi.org/10.1088/1755-1315/562/1/012015>.
- Wikantiyoso, Respati, Pindo Tutuko, Tonny Suhartono, Aditya Galih Sulaksono, and Adisti Safrilia. 2020. 'Green City MIS as a Sustainable Urban GOS Provision Control Implementation Model'. *International Review for Spatial Planning and Sustainable Development* 8 (1): 160–72. https://doi.org/10.14246/irspda.8.1_160.
- Xu, Dandan. 2014. 'Compare NDVI Extracted from Landsat 8 Imagery with That from Landsat 7 Imagery'. *American Journal of Remote Sensing* 2 (2): 10. <https://doi.org/10.11648/j.a.jrs.20140202.11>.
- Yunhao, Chen, Shi Peijun, Li Xiaobing, Chen Jin, and Li Jing. 2006. 'A Combined Approach for Estimating Vegetation Cover in Urban/Suburban Environments from Remotely Sensed Data'. *Computers & Geosciences* 32 (9): 1299–1309. <https://doi.org/10.1016/j.cageo.2005.11.011>.
- Zaitunah, A, Samsuri, A G Ahmad, and R A Safitri. 2018. 'Normalized Difference Vegetation Index (Ndvi) Analysis for Land Cover Types Using Landsat 8 Oli in Besitang Watershed, Indonesia'. *IOP Conference Series: Earth and Environmental Science* 126 (March): 012112. <https://doi.org/10.1088/1755-1315/126/1/012112>.

Detection of potential green open space area using landsat 8 satellite imagery

ORIGINALITY REPORT

21 %
SIMILARITY INDEX

14 %
INTERNET SOURCES

16 %
PUBLICATIONS

15 %
STUDENT PAPERS

PRIMARY SOURCES

1 Submitted to Universitas Riau
Student Paper 1 %

2 bdm.unb.br
Internet Source 1 %

3 Submitted to Hong Kong Baptist University
Student Paper 1 %

4 Hamere Yohannes, Teshome Soromessa, Mekuria Argaw, Ashraf Dewan. "Changes in landscape composition and configuration in the Beressa watershed, Blue Nile basin of Ethiopian Highlands: historical and future exploration", Heliyon, 2020
Publication 1 %

5 Submitted to UPN Veteran Jawa Timur
Student Paper 1 %

6 ejournal.undip.ac.id
Internet Source 1 %

7 Hong Tao, Manqi Li, Ming Wang, Guonian Lü. "Genetic algorithm-based method for forest

type classification using multi-temporal NDVI from Landsat TM imagery", Annals of GIS, 2018

Publication

8	scholar.unand.ac.id Internet Source	1 %
9	Submitted to Salem State University Student Paper	1 %
10	Y Basuki, Widjonarko. "Hydrometeorological hazard prediction in the kuto bodri river region central java based on normalized difference vegetation index (ndvi) analysis", IOP Conference Series: Earth and Environmental Science, 2021 Publication	1 %
11	journal.unj.ac.id Internet Source	1 %
12	zookeys.pensoft.net Internet Source	1 %
13	Submitted to Maulana Azad National Institute of Technology Bhopal Student Paper	1 %
14	Submitted to Universitas Amikom Student Paper	1 %
15	Submitted to SDM Universitas Gadjah Mada Student Paper	1 %

16	Submitted to Universitas Hasanuddin Student Paper	1 %
17	repo.undiksha.ac.id Internet Source	1 %
18	documents1.worldbank.org Internet Source	1 %
19	www.geosystems.de Internet Source	1 %
20	abis.kastamonu.edu.tr Internet Source	1 %
21	enveurope.springeropen.com Internet Source	1 %
22	Xiaoxian Ju. "Application of Big Data Technology to Promote Agricultural Structure Adjustment and High-Quality Development of Modern Agriculture", Computational Intelligence and Neuroscience, 2022 Publication	1 %
23	annalsofrscb.ro Internet Source	1 %
24	media.neliti.com Internet Source	1 %
25	ojs.unida.ac.id Internet Source	1 %

26

api.research-repository.uwa.edu.au

Internet Source

1 %

27

ejournal2.undip.ac.id

Internet Source

1 %

Exclude quotes Off

Exclude matches < 1%

Exclude bibliography On

Detection of potential green open space area using landsat 8 satellite imagery

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6
